

**B.TECH.**  
**THEORY EXAMINATION (SEM-II) 2016-17**  
**ENGINEERING MATHEMATICS - II**

Time : 3 Hours

Max. Marks : 100

Note : Be precise in your answer. In case of numerical problem assume data wherever not provided.

**SECTION – A**

1. Explain the following: **10 x 2 = 20**

- (a) Show that the differential equation  $y dx - 2x dy = 0$  represents a family of parabolas.
- (b) Classify the partial differential equation
 
$$(1 - x^2) \frac{\partial^2 z}{\partial x^2} - 2xy \frac{\partial^2 z}{\partial y \partial x} + (1 - y^2) \frac{\partial^2 z}{\partial y^2} = 2z$$
- (c) Find the particular integral of  $(D - \alpha)^2 y = e^{\alpha x} f''(x)$ .
- (d) Write the Dirichlet's conditions for Fourier series.
- (e) Prove that  $J'_0(x) = -J_1(x)$ .
- (f) Prove that  $L [e^{at} f(t)] = F(s - a)$
- (g) Find the Laplace transform of  $f(t) = \frac{\sin at}{t}$ .
- (h) Write one and two dimensional wave equations.
- (i) Find the constant term when  $f(x) = |x|$  is expanded in Fourier series in the interval  $(-2, 2)$ .
- (j) Write the generating function for Legendre polynomial  $P_n(x)$ .

**SECTION – B**

2. Attempt any five of the following questions: **5 x 10 = 50**

- (a) Solve the differential equation
 
$$(D^2 + 2D + 2)y = e^{-x} \sec^3 x, \quad \text{where } D = \frac{d}{dx}$$
- (b) Prove that  $(n + 1)P_{n+1}(x) = (2n + 1)xP_n(x) - nP_{n-1}(x)$ , where  $P_n(x)$  is the Legendre's function.
- (c) Find the series solution of the differential equation
 
$$2x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - (x + 1)y = 0.$$
- (d) Using Laplace transform, solve the differential equation
 
$$\frac{d^2 y}{dt^2} + 9y = \cos 2t; \quad y(0) = 1, \quad y\left(\frac{\pi}{2}\right) = -1.$$
- (e) Obtain the Fourier series of the function,
 
$$f(t) = t, \quad -\pi < t < 0$$

$$= -t, \quad 0 < t < \pi.$$
 Hence, deduce that  $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$
- (f) Solve  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$  under the conditions  $u(0, y) = 0$ ,
 
$$u(l, y) = 0, u(x, 0) = 0 \text{ and } u(x, a) = \sin \frac{n\pi x}{l}.$$
- (g) Solve the partial differential equation:
 
$$(D^3 - 4D^2 D' + 5D D'^2 - 2D'^3)z = e^{y+2x} + \sqrt{y+x}$$
- (h) Using convolution theorem find  $L^{-1} \left[ \frac{1}{(s+1)(s^2+1)} \right]$

## SECTION – C

2 x 15 = 30

Attempt any two of the following questions:

3. (a) Solve the differential equation  $(D^2 - 2D + 1)y = e^x \sin x$
- (b) Solve the equation by Laplace transform method:  

$$\frac{dy}{dt} + 2y + \int_0^t y dt = \sin t, \quad y(0) = 1.$$
- (c) Solve the partial differential equation  

$$(y^2 + z^2)p - xyq + zx = 0, \text{ where } p = \frac{\partial z}{\partial x} \text{ \& } q = \frac{\partial z}{\partial y}$$
4. (a) Find the Laplace transform of  $\frac{\cos at - \cos bt}{t}$ .
- (b) Express  $f(x) = 4x^3 - 2x^2 - 3x + 8$  in terms of Legendre's polynomial.
- (c) Expand  $f(x) = 2x - 1$  as a cosine series in  $0 < x < 2$ .
5. (a) Show that  $J_3(x) = \left(\frac{8}{x^2} - 1\right)J_1(x) - \frac{4}{x}J_0(x)$ .
- (b) Solve the  $2 \frac{\partial z}{\partial x} + 3 \frac{\partial z}{\partial y} + 5z = 0; z(0, y) = 2e^{-y}$  by the method of separation of variables.
- (c) A tightly stretched string with fixed end  $x = 0$  and  $x = l$  is initially in a position given by  $y = a \sin \frac{\pi x}{l}$ . If it is released from rest from this position, find the displacement  $y(x, t)$ .